



Sick Child Policy

FFCC SICK Child Policy

FFCC realizes that most of you are working parents and we do try to accommodate you as much as possible. However, for the health and wellbeing of the children in our care and our staff we feel it is vital to maintain a strict wellness policy. We ask for parents to assist by keeping sick children at home. We do take into consideration your physician's recommendations nevertheless it is up to our discretion when your child may return to the center.

Fairlane Family Child Care reserves the right to temporarily deny any child admittance to the center for reasons of obvious illness, or to request early departure should symptoms become apparent during the course of the day. Any child who seems unable to participate in our program for any reason will be sent home. This is to ensure the continued good health of everyone at the center.

When a child may not attend school:

- **Fever:** Children will be sent home if their temperature is 100.0 or higher and must stay home the next day for observation. Children must be free of fever (any temperature above 98.6 degrees) for at least 24 hours without the use of fever reducing medication. The same policy applies if your child develops a fever at home. They must be fever free (any temperature above 98.6 degrees) for at least 24 hours without the use of fever reducing medication.
- **Rash:** Any rash other than a common diaper rash or skin irritation will require that child to be sent home for an evaluation and diagnosis from their doctor in writing of exactly what it is. They may return to school based on that written doctor's evaluation, and clearance that it is not contagious.
- **Conjunctivitis (pink eye):** Children will be sent home if there appears to be an unusual amount of discharge from or irritation to their eye(s) and must stay home the next day for observation. Before returning to school they will need an evaluation and diagnosis from their doctor in writing of exactly what it is.
- If the diagnosis is **BACTERIAL CONJUNCTIVITIS** children must have received at least 24 hours of treatment.
- If the diagnosis is **VIRAL CONJUNCTIVITIS** your child may return **AS LONG AS THERE IS NO DISCHARGE**.
- If in fact they do not have "pink eye" we need a doctor's note with a diagnosis and a clearance that it is not contagious.
- **Thick White, Green or Yellow Discharge:** Children will be sent home if they appear to have any thick white, green or yellow discharge. This is often indicative of an infection and they must stay

home the next day for observation. Before returning to school they will need an evaluation and diagnosis from their doctor in writing and at least 24 hours of treatment. If in fact they do not have

an infection we need a doctor's note with a diagnosis of exactly what it is with a clearance that it is nothing contagious. The Department of Health and most doctors are of the opinion that once on antibiotics for 24 hours, the discharge is no longer contagious even though it may persist for up to two weeks.

- **Diarrhea:** Children will be sent home if they have three or more loose bowel movements in one day and must stay home the next day for observation. Before returning to school (after the day of observation) children must be free from diarrhea for 24 hours with at least 1 regular bowel movement. If your child has one or more loose bowel movements on their first day back they will again be sent home.
- **Vomiting:** Children will be sent home if they vomit and must stay home the next day for observation. Before returning to school (after the day of observation) children must symptom free with no vomiting for at least 24 hours.
- **Persistent Hacking Cough:** Children will be sent home if they have a persistent hacking cough and must stay home the next day for observation. Before returning to school they will need an evaluation and diagnosis from their doctor in writing and at least 24 hours of treatment. If in fact child do not require treatment, we need a doctor's note that it is not contagious.
- **Lice:** Children will not be readmitted until 24 hours after treatment and must be nit free. We will make an evaluation and determine if the child can be readmitted.

Common Cold Policy

Children suffering from a common cold will be assessed on an individual basis.

Factors of consideration include the developmental level of your child in congruence with our ability to limit the spread of germs.

The younger your child, the more difficult it is to keep the spread of germs down. For example: hand to face contact, mouthing of toys, uncontrolled nasal discharge, uncovered sneezing and coughing etc.

Medication Administration Policy

Medications both prescription and over the counter are rarely given at school; the only exceptions involve special or serious problems where it deemed absolutely necessary by the physician that the medication be given during school hours. The parent is urged, with the help of your child's physician, to work out a schedule of giving medication at home, outside school hours whenever possible.

IF MEDICATION IS TO BE ADMINISTERED AT SCHOOL, ALL OF THE FOLLOWING CONDITIONS MUST BE MET

1. A signed request from a licensed physician/dentist specifying the condition for which the medication is to be given, the name, dosage, route, side effect and specific instructions for emergency treatment must be on file at school. School staff is not authorized to determine when an "as needed" medication is to be given. Specific instructions are necessary.
2. A signed request from the parent/guardian must be on file at school.
3. Medication must be in your child's original, labeled pharmacy container written in English.
4. All liquid medication must be accompanied by an appropriate measuring device.
5. A separate form is required for each medication.
6. We do not administer over the counter medicine which includes any medicine for allergies, fever, cold, cough or any pain medications without a doctor's note and signed forms along with instructions.

Keep Me Home If...



Seattle King County Department of Public Health

When Your Child is Sick:

1. Have plans for back up child care.
2. Tell your caregiver what is wrong with your child, even if your child stays home.

I NEED TO STAY AT HOME IF...

FEVER	VOMITING	DIARRHEA	RASH	HEAD LICE	EYE INFECTION	HOSPITAL STAY AND/OR ER VISIT
Temperature of 100.4 or higher	Within the past 24 hours	Within the past 24 hours	Body rash with itching or fever	Itchy head, active head lice	Redness, itching, and/or "crusty" drainage from eye	Hospital stay and/or ER visit

I AM READY TO GO BACK TO WORK OR SCHOOL WHEN I AM...

Fever free for 24 hours without the use of fever reducing medication i.e. Tylenol, Motrin	Free from vomiting for at least 2 solid meals	Free from diarrhea for at least 24 hours	Free from rash itching, or fever. I have been evaluated by my doctor if needed.	Treated with appropriate lice treatment at home and proof is provided to nurse.	Evaluated by my doctor and have note to return to school.	Released by medical provider to return to work or school.
---	---	--	---	---	---	---

Coronavirus Disease 2019 (COVID-19) Outbreak Update

Maryland state and Howard County government have declared “the state of emergency” due to Coronavirus Disease 2019 (COVID-19) Outbreak in Maryland state and in Howard County. All county schools are closed for two weeks starting from March 16th 2020. We are closely monitoring the situation and may close the center if outbreak reaches to our communities or if it poses any threats to children in our care or any of staff members to keep everyone safe. You are subject to pay for the closing period in case of state of emergency or any reasons beyond our control to keep the spot for your child. We will send an email with dates to all parents if we decide to close to help slowing down the outbreak as per county and state guidelines.

If you have travelled out of country/state or attended large gatherings or used public transportation, please consider keeping child home for 48 hours and monitor before returning to the center,

Meanwhile, please see following guidelines from CDC for coronavirus (COVID-19).

For more information: www.cdc.gov/COVID19

What to do if you are sick with coronavirus disease 2019 (COVID-19)

If you are sick with COVID-19 or suspect you are infected with the virus that causes COVID-19, follow the steps below to help prevent the disease from spreading to people in your home and community.

Stay home except to get medical care

You should restrict activities outside your home, except for getting medical care. Do not go to work, school, or public areas. Avoid using public transportation, ride-sharing, or taxis.

Separate yourself from other people and animals in your home

People: As much as possible, you should stay in a specific room and away from other people in your home. Also, you should use a separate bathroom, if available.

Animals: Do not handle pets or other animals while sick. See [COVID-19 and Animals](#) for more information.

Call ahead before visiting your doctor

If you have a medical appointment, call the healthcare provider and tell them that you have or may have COVID-19. This will help the healthcare provider's office take steps to keep other people from getting infected or exposed.

Wear a facemask

You should wear a facemask when you are around other people (e.g., sharing a room or vehicle) or pets and before you enter a healthcare provider's office. If you are not able to wear a facemask (for example, because it causes trouble breathing), then people who live with you should not stay in the same room with you, or they should wear a facemask if they enter your room.

Cover your coughs and sneezes

Cover your mouth and nose with a tissue when you cough or sneeze. Throw used tissues in a lined trash can; immediately wash your hands with soap and water for at least 20 seconds or clean your hands with an alcohol-based hand sanitizer that contains at least 60-95% alcohol covering all surfaces of your hands and rubbing them together until they feel dry. Soap and water should be used preferentially if hands are visibly dirty.

Avoid sharing personal household items

You should not share dishes, drinking glasses, cups, eating utensils, towels, or bedding with other people or pets in your home. After using these items, they should be washed thoroughly with soap and water.

Clean your hands often

Wash your hands often with soap and water for at least 20 seconds. If soap and water are not available, clean your hands with an alcohol-based hand sanitizer that contains at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Soap and water should be used preferentially if hands are visibly dirty. Avoid touching your eyes, nose, and mouth with unwashed hands.

Clean all “high-touch” surfaces every day

High touch surfaces include counters, tabletops, doorknobs, bathroom fixtures, toilets, phones, keyboards, tablets, and bedside tables. Also, clean any surfaces that may have blood, stool, or body fluids on them. Use a household cleaning spray or wipe, according to the label instructions. Labels contain instructions for safe and effective use of the cleaning product including precautions you should take when applying the product, such as wearing gloves and making sure you have good ventilation during use of the product.

Monitor your symptoms

Seek prompt medical attention if your illness is worsening (e.g., difficulty breathing). **Before** seeking care, call your healthcare provider and tell them that you have, or are being evaluated for, COVID-19. Put on a facemask before you enter the facility. These steps will help the healthcare provider's office to keep other people in the office or waiting room from getting infected or exposed.

Ask your healthcare provider to call the local or state health department. Persons who are placed under active monitoring or facilitated self-monitoring should follow instructions provided by their local health department or occupational health professionals, as appropriate.

If you have a medical emergency and need to call 911, notify the dispatch personnel that you have, or are being evaluated for COVID-19. If possible, put on a facemask before emergency medical services arrive.

Discontinuing home isolation

Patients with confirmed COVID-19 should remain under home isolation precautions until the risk of secondary transmission to others is thought to be low. The decision to discontinue home isolation precautions should be made on a case-by-case basis, in consultation with healthcare providers and state and local health departments.



CS 134837-0 03/05/2020

For more information: www.cdc.gov/COVID19